milies
uth

Table of Contents

PREFACE..... 1 ;

Preface

Each day, the safety and well-being of children across the Nation are threatened by child abuse and neglect. Many of these children live in homes where substance use disorders (substance abuse and dependence) create additional and compounding problems. e child welfare and alcohol and drug abuse treatment “elds are working to “nd e ective ways to serve families where this overlap occurs. Intervening e ectively in the lives of these children and their families is not the responsibility of a single agency or professional group, but rather it is a shared community concern.

Since the late 1970s, the *Child Abuse and Neglect User Manual Series* has provided guidance on child protection to hundreds of thousands of multidisciplinary professionals and concerned community members. e *User Manual Series* o ers a foundation for understanding child maltreatment and the roles and responsibilities of various practitioners in its prevention, identi“cation, investigation, and treatment. rough the years, the manuals have served as valuable resources for building knowledge, promoting e ective practices, and enhancing community collaboration.

A number of changes have occurred that dramatically a ect each community•s response to child maltreatment since the last update of the *User Manual Series* in the early 1990s. e changing landscape includes increased recognition of the complexity of issues facing children and families, new legislation, practice innovations, and systems reform e orts. Significant advances in research have helped shape new directions for interventions, while ongoing evaluations show what works.Ž

e O ce on Child Abuse and Neglect within the Children•s Bureau of the Administration for Children and Families, U.S. Department of Health and Human Services, has developed this third edition of the *User Manual Series* to re”ect increased knowledge and the evolving state of practice on child protection. e updated and new manuals are comprehensive in scope, while succinct in presentation and easy to follow, and they address trends and concerns relevant to today•s professional.

is manual, *Protecting Children in Families A ected by Substance Use Disorders*, provides a basis for understanding parental substance use disorder and its relationship to child maltreatment. e manual is intended to help child protective services (CPS) caseworkers identify when drug and alcohol use by a parent or other caregiver is a factor in child welfare cases and work more e ectively to meet the needs of all family members involved. e manual encourages enhanced collaboration between CPS and alcohol and drug abuse treatment agencies. While CPS caseworkers are the primary audience, alcohol and drug abuse treatment providers may “nd the manual helpful in building their understanding of child protection issues among the families with whom they work. It also may be useful to other professionals,

Acknowledgments

Contributing Authors

Kathleen Feidler, M.S.W., a consultant with ICF International, has more than 15 years of experience in the evaluation of substance use disorder programs. She also has researched and written several articles and papers about the impact of substance use disorder in child welfare cases.

Karen Mooney, L.C.S.W., C.A.C. III, is the women•s treatment coordinator for the Alcohol and Drug Abuse Division of the Colorado Department of Human Services. Formerly, she was a child protective services caseworker for 7 years, during which she specialized in screening, assessing, and engaging parents of children involved in the child welfare system in substance use disorder and addiction treatment. Since that time, she has served as a child protection and substance use disorder treatment consultant to several national organizations and universities.

Mary Nakashian, M.A., is a consultant specializing in public policy and management, training, and technical

Additional Acknowledgments

e third edition of the

Purpose and Overview

The relationship between substance use disorders (SUDs) and child maltreatment is compelling and undeniable. More than eight million children in the United States live with at least one parent who abused or was dependent on alcohol or an illicit drug during the past year.[1] These children face a heightened risk of maltreatment.[2] One study, for example, showed that children of parents with SUDs are nearly three times more likely to be abused and more than four times more likely to be neglected than children of parents who do not abuse substances.[3]

According to the National Child Abuse and Neglect Data System (NCANDS), in 2006, an estimated 3.3 million referrals were made to child protective services (CPS), representing 6 million children. From this population, approximately 905,000 children were found to be victims of child abuse or neglect. Of the maltreated children, 66.3 percent were neglected (including medical neglect), 16.0 percent physically abused, 8.8 percent sexually abused, and 6.6 percent psychologically maltreated. Additionally, 15.1 percent of victims were associated with •otherŽ types of maltreatment, such as abandonment or congenital drug addiction. A child could be identi“ed as a victim of more than one type of maltreatment.[4] Additionally, while estimates vary, most studies suggest that parental

7KLV SDJH LV LQWHQWLRQDOO\ OHIW EODQ

to buying stolen drugs. Exhibit 2-1 provides key statistics for commonly abused substances.

Other substances may not have medicinal qualities but can a ect users psychologically and physically or lower inhibitions and impair judgment if misused. For instance, some individuals drink alcohol at social gatherings to feel more comfortable talking and relating to others. Being of legal age and drinking alcohol is a commonly accepted practice in the United States. Of course, alcohol often can be misused and can negatively a ect events ranging from traffic safety to the ability to care adequately for children.

Exhibit 2-1
Selected Drug Statistics from the National Survey on Drug Use and Health (NSDUH)*

- An estimated 19.9 million Americans, or 8.0 percent of the population aged 12 or older, were current illicit drug users in 2007.** (is "gure re"ects use of the following drugs: marijuana, cocaine, heroin, hallucinogens, and inhalants and the nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, and sedatives.)

- e estimated number of Americans who were current users of the following drugs in 2007:

  - Marijuana: 14,448,000
  - Cocaine: 2,075,000 (including 610,000 users of crack)
  - Hallucinogens: 996,000 (including 503,000 users of Ecstasy)
  - Inhalants: 616,000
  - Heroin: 338,000

- In 2007, approximately 6.9 million people aged 12 or older (2.8 percent of the population) were current users of prescription-type psychotherapeutic drugs taken nonmedically, including pain relievers, tranquilizers, stimulants, and sedatives. is includes 529,000 individuals who were current users of methamphetamine, which can be manufactured illegally using existing prescription drugs.

- An estimated 22.3 million Americans aged 12 or older in 2006 (9.0 percent of the population) were classi"ed with substance abuse or dependence. Of these:

  - 3.2 million abused or were dependent on both alcohol and illicit drugs;
  - 3.7 million abused or were dependent on illicit drugs but not alcohol;
  - 15.5 million abused or were dependent on alcohol but not illicit drugs.[10]

* ese statistics are drawn from the 2007 NSDUH, an annual survey of the civilian, noninstitutionalized population of the United States aged 12 or older. To see the full results of the most recent survey, visit the website of the U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, O ce of Applied Studies: http://www.oas.samhsa.gov/nsduhLatest.htm.

** •Current usersŽ re"ect persons who used the speci"ed drug during the month prior to the NSDUH interview.

Quick Facts on Alcohol Use

- Slightly more than half of Americans aged 12 or older, or approximately 127 million people, reported being current drinkers of alcohol in the 2007 NSDUH. (Current drinkers were de“ned as having had at least one drink in the 30 days prior to the survey.)
- An estimated 17 million people (6.9 percent of the population) were heavy drinkers. (Heavy drinking was de“ned as having “ve or more drinks on the same occasion on at least 5 di erent days in the past 30 days.)
- Among pregnant women aged 15 to 44, an estimated 11.6 percent reported current alcohol use, and 3.7 percent reported binge drinking. (Binge drinking was de“ned as having “ve or more drinks on the same occasion on at least 1 day in the past 30 days.)
- Excessive alcohol use is the third leading lifestyle cause of death in the United States and was determined to be a key factor in approximately 79,000 deaths annually from 2001…2005.
- e U.S. Dietary Guidelines for Americans recommends no more than one drink per day for adult women and no more than two drinks per day for adult men. It also lists several types of individuals„ including children, adolescents, and pregnant women„who should avoid alcohol completely.[11]

For more information on commonly abused substances, see Appendix E, *Commonly Abused Substances*.

With respect to child protection, substance use becomes problematic when it contributes to the harm of children. is can be di cult for CPS caseworkers to identify because the distinction between •normalŽ alcohol use and problematic use may be blurred and subject to interpretation. (See Chapter 4, *In-home Examination, Screening, and Assessment of Substance Use Disorders*, for more information about identifying SUDs.)

Characteristics of Addiction

Knowing the characteristics of addiction can help inform e ective intervention and practice with individuals su ering from SUDs. Characteristics include:

- \ Progressive Nature. A central feature of addiction is a progressive use of a substance, whether alcohol, prescription medications, or illegal drugs. e physical, emotional, and social problems that arise from addiction typically continue to worsen unless the SUDs are treated successfully. If left untreated, addiction can cause premature death through overdose; through organic complications involving the brain, liver, heart, and many other organs; and by contributing to motor vehicle crashes, homicide, suicide, and other traumatic events.
- \ Denial and Concealment. Addiction can be di cult to identify, even for individuals experiencing it. People who are addicted to a substance often engage in elaborate strategies to conceal the amount being consumed and the degree to which the substance is ista ing their lives. Another dimension of addiction is that individuals who su er from it often do not

perceive that their pattern of drinking or drug use creates or contributes to their problems. Additionally, the use of substances may cause their memory or perception of events or of what they have said or done. is lack of recognition commonly is identi“ed as denial.

- Chronic Disease. e National Institute on Drug Abuse has de“ned addiction as a chronic disease, like heart disease, hypertension, and diabetes. Studies have shown alcohol and drug abuse treatment is about as e ective as treatments required for these other chronic diseases. Lifetime management of chronic diseases in all cases requires individuals to change their habits and activities and to take precautions that prevent them from relapsing or worsening their condition.

- Lapses and Relapses. Lapses and relapses are common features of addiction. A lapse is a period of substance use after the individual has been clean and sober for some length of time. A relapse is not only using the substance again, but also returning to the problem behaviors associated with it.[12]

Addiction is di cult to deal with; many individuals lapse or relapse one or more times before being able to remain abstinent. If lapses or relapses occur, they do not necessarily mean that treatment has failed. ey can point the way toward needed improvements in how those individuals are approaching recovery. Most individuals who have lapsed or relapsed can identify, prior to the lapse or relapse, certain situations, thoughts, or behaviors that contributed to the use of the substance.

Why Some People Become Addicted

Many theories and explanations have been proposed to describe the reasons why some individuals become addicted to substances and others do not. Research on the causes of addiction is not conclusive, and multiple factors may contribute to it. Early explanations for addiction included moral weakness, insanity, demonic possession, and character pathology.[13] ese explanations, combined with the problematic behaviors that sometimes accompany addiction, have created a serious stigma. Recent research, however, indicates that substance addiction is a brain disease that changes its structure and functioning, which in turn a ects an individual•s behaviors. Although the initial use of a substance may be voluntary, a person•s ability to control future use may be seriously impaired by changes in the brain caused by prior use.[14]

Some research, including adoption and twins studies, has demonstrated a biological and genetic predisposition to addiction, with scientists estimating that genetic factors account for 40…60 percent of an individual•s risk of addiction.[15] ese studies suggest that an individual•s genes play a role in vulnerability to addiction. For example, one study found that children whose parents are addicted to drugs or alcohol are three times more likely to develop an SUD later in life than children whose parents are not addicted.[16] Other research emphasizes a social factor to explain that addictions appear to •run in the family.Ž ese studies suggest that children who grow up in families with SUDs may model their adult behavior on what they have seen and known in their familial experience.[17] Risk for addiction can also be a ected by gender, ethnicity, developmental stage, and social environment.[18] In other words, both nature and nurture contribute to a person•s vulnerability or resistance to substance abuse.

Many self-help groups, such as 12-step programs, consider addiction a progressive illness that is physical, spiritual, and emotional in nature. ey believe that individuals who are addicted must admit that they are powerless over the substance; that is, they are unable to resolve the problem on their own and must seek help outside themselves.[19]

Psychophysical withdrawal occurs when individuals experience physical symptoms that result from

of the batterers• violence and interfere with domestic violence interventions.[28] ey also contribute to the increased severity of injuries among victims.[29]

Poverty

SUDs cross all socioeconomic lines, but studies show that there is a relationship between poverty and substance abuse.[33] People living in poverty sometimes turn to substances for relief from the anxiety and the stress associated with economic insecurity. Of course, spending money on alcohol or drugs often only contributes to economic problems. Dealing illegal drugs is viewed by some as a source of income and a means of escaping poverty. Unfortunately, some individuals su ering from economic hardship feel that they have little to lose if they get involved in drugs, no matter what the e ects are on themselves or their families.

Parents who are distracted by their nancial problems may have less energy and attention for parenting. In some homes, the psychological distress of poverty may be directed toward the children. Research has indicated a strong association between child maltreatment, particularly neglect, and poverty.[34] CPS case plans invariably need to address issues related to poverty and establish service plans for families.

Homelessness

In some cases, extreme poverty and other factors may lead to homelessness. Homeless people typically experience several overlapping challenges, including SUDs, mental illnesses, and a variety of physical health problems. Parents with children account for approximately 11 percent of the homeless population, and this number appears to be growing.[35]

Crime ;

Crime has a strong association with drug use. In the most recent study of its kind, more than three out of every four State, Federal, or local jail inmates previously were involved seriously with drugs or alcohol in some way (e.g., convicted of a drug- or alcohol-related crime, used illicit substances regularly, were under the in”uence of alcohol or drugs when they committed a crime).[rÍ üÎBnØ¢ Ø¬¬„„H n‰ ÊÈÒoæ‘Ì”ytv¦Ìª îiú•® Ä—Þîº ◦

childhood development. It is intended to help child

directly a ects the fetus.[58] A variety of birth defects to the major organs and the central nervous system, which are permanent, can occur due to alcohol use during pregnancy, though the risk of harm decreases if the pregnant woman stops drinking completely.[59] Collectively, these defects are called Fetal Alcohol Syndrome (FAS). FAS is one of the most commonly known birth defects related to prenatal drug exposure. Children with FAS may exhibit:

- Growth de“ciencies, both prenatally and after birth
- Problems with central nervous system functioning
- IQs in the mild to severely retarded range
- Small eye openings and poor development of the optic nerve
- A small head and brain
- Joint, limb, ear, and heart malformations.

Alcohol-Related Neurodevelopmental Disorder (ARND) and Alcohol-Related Birth Defects (ARBD) are similar to FAS. Once known as Fetal Alcohol Effects, ARND and ARBD are terms adopted in 1996 by the National Academy of Sciences• Institute of Medicine. ARND and ARBD encompass the functional and physiological problems associated with prenatal alcohol exposure, but are less severe than FAS. Children with ARND can experience functional or mental impairments as a result of prenatal alcohol exposure, and children with ARBD can have malformations in the skeletal and major organ systems. Not all children who are exposed prenatally to alcohol develop FAS, ARND, or ARBD, but for those who do, these e ects continue throughout their lives and at all the stages of development, although they are likely to present themselves di erently at each developmental stage. Exhibit 3-1 compares typical childhood behavior at each developmental stage with behaviors and characteristics associated with FAS, ARND, and ARBD.

Exhibit 3-1
Childhood Behavior and Characteristics Associated with FAS, ARND, and ARBD[60]

Developmental Stage	Typical Behaviors or Characteristics	FAS/ARND/ARBD Behaviors or Characteristics
Infants	• Develop mental and physical skills • Bond with caretakers	• Problems with spatial and depth perception, muscle coordination and development, facility with speech, and processing information • Attention de“cit disorder • Inability to focus • Possible attachment disorders
Toddlers	• Develop sense of self • Assert independence by saying •noŽ	• Di culty exercising self-control, which leads to self> doubt and feelings of inadequacy
5…7 year olds	• Try new things • Meet or exceed academic standards • Learn new social skills	• Overwhelmed with new situations and interactions with other children • Inability to pick up social skills by observation • Problems meeting academic standards
8…12 year olds	• Increased in”uence of peers • Games become important method of bonding and developing interpersonal skills	• Di culty remembering rules of games • Lack of remorse in breaking rules • Become depressed and exhibit other behavior problems
Teenagers	• Continued detachment from parents • Development of individual identity • Learn to identify with larger community	• May lack skills to become good community members • Become socially isolated • May “nd their way to peer groups that engage in high> risk behaviors • May withdraw altogether from groups

The Impact on Childhood Development

Exposure to parental SUDs during childhood also can have dire consequences for children. Compared to children of parents who do not abuse alcohol or drugs, children of parents who do, and who also are in the child welfare system, are more likely to experience physical, intellectual, social, and emotional problems. Among the di culties in providing services to these children is that problems a cted or compounded by their parents• SUDs might not emerge until later in their lives.[68]

is section summarizes some of the consequences of SUDs on childhood development, including a disruption of the bonding process; emotional, academic, and developmental problems; lack of supervision; parenti“cation; social stigma; and adolescent substance use and delinquency.

Disruption of the Bonding Process

When mothers or fathers abuse substances after delivery, their ability to bond with their child„so important during the early stages of life„may be weakened. In order for an attachment to form, it is necessary that caregivers pay attention to and notice their children•s attempts to communicate. Parents who use marijuana, for example, may have di culty picking up their babieight not eihEs obecase marijuana,durll senspnseq tim and daltrs paeocetion . ents senpete.dy smisstheir babieight not eihEs ,the bobieig velnual,ly wstopproviding she m ensltyis ents

theinhave di culty , tapro
aenltion;ship16(v.)]TJ EMC /P <</MCID 45>>BDC 0.12062Tw 0 -1.189 Td [(cN30(reglcted ohildr)10(en iho al)
wasof lthers ht not eiaotionas Inpoaiendeocial scogniion;
pcn alar obpaeowdaynpcn alar obpae2 Tw T* [(1asof lthers tfuls)isTJ 0 /Reference <</MCID 11 >>BDC 22T0

In-home Examination, Screening, and Assessment of Substance Use Disorders

In This Chapter

- In-home examination
- Screening
- Assessment

The previous chapter described some of the effects of parental substance use disorders (SUDs) on children, but how does a child protective services (CPS) worker determine if SUDs exist in the family? This chapter discusses in-home examinations, SUD screening instruments, and SUD assessments, including their methods, benefits, and limitations, and how caseworkers can incorporate them into their practice.

In-home Examination

An in-home examination includes observations by the CPS caseworker of the people and the environment in a home. When visiting a home as part of an investigation, the caseworker should check for the following indicators of possible SUDs:

- A report of substance use is included in the CPS call or report
- Drug paraphernalia (e.g., a syringe kit, charred spoons, a large number of liquor or beer bottles)
- The scent of alcohol or drugs
- A child or other family member reports alcohol or drug use by a parent
- A parent appears to be under the influence of a substance, admits to having an SUD, or shows other signs of addiction or abuse (e.g., needle marks).

This list can be used pre- or post-screening and can be incorporated into every home visit.[77]

Screening

Screening is the use of a simple, and usually brief, set of questions that have been validated (i.e., tested to show that they accurately indicate the presence of an SUD). Results are easy to interpret. Generally, individuals who are not trained SUD treatment providers use the instruments.

The goal of screening is to determine whether a family member requires further evaluation for SUDs.[78] CPS caseworkers can use screening as a part of their standard home visits or family assessments. This section describes the importance of screening, sample screening instruments, benefits and limitations of screening instruments, and what to do when an instrument indicates an individual may have an SUD.

Signs of Methamphetamine Use and Manufacture

With the increased use of methamphetamine, “rst responders are now more likely to work with clients who are users or manufacturers of this drug. e following information can assist them in identifying methamphetamine use or manufacturing.

Signs of possible methamphetamine use include:

- Increased breathing and pulse rate
- Sweating
- Rapid/pressured speech
- Euphoria (an exaggerated feeling of well-being)
- Hyperactivity
- Dry mouth
- Tremors (shaking hands)
- Dilated pupils
- Lack of appetite
- Insomnia or lack of sleep
- Bruxism (teeth-grinding)
- Depression
- Irritability, suspiciousness, paranoia
- Visual and auditory hallucinations
- Formication (the sensation of bugs crawling on the skin)
- e presence of white powder, straws, or injection equipment.

Signs that methamphetamine is possibly being manufactured in a home include:

- Laboratory equipment (e.g., ”asks, rubber tubing, beakers, large amounts of coffee filters)
- Large quantities of pills containing ephedrine or pseudoephedrine (e.g., certain cold medicines)
- A chemical odor
- Chemicals not commonly found in a home (e.g., red phosphorous, acetone, liquid ephedrine, ether, iodine, phenylacetone [P2P])

are more apparent. e opposite also can be true: an SUD may mask other problems, such as domestic violence or disabilities. Moreover, addiction often is characterized by denial, and family members may not recognize that they have an SUD. For example, they may feel that their drinking is within an acceptable range or that their marijuana use is not problematic.

Screening is just one of many approaches used to identify SUDs. It is not completely accurate nor will it work all the time (i.e., not all positive responses will demonstrate an SUD, and not all negative responses will rule out a disorder). CPS caseworkers should also rely on additional techniques, such as observation, medical histories, reports from family members or friends, or arrest records. Nevertheless, using a brief screening instrument takes little time and provides an objective method for caseworkers to use in opening a discussion about sensitive issues. In this way, screening becomes part of a continuum of activities aimed at addressing families• problems with SUDs.

Sample Screening Instruments

Ideally, screening instruments used by CPS < caseworkers should be brief, easily administered, <

Exhibit 4-1
Sample Screening Instruments for Substance Use Disorders

e CAGE Questionnaire (amended for drug use)

C Have you ever felt the need to Cut down on your drinking or drug use?

A Have you ever felt Annoyed by people criticizing your drinking or drug use?

G Have you ever felt bad or Guilty about your drinking or drug use?

E Have you ever had a drink or used a drug “rst thing in the morning to steady your nerves or get rid of a hangover? (Eye-opener)

Scoring: If the answer is •yesŽ to one or more questions, the responder should receive a formal alcohol and drug assessment. Answering •yesŽ to one or two questions may indicate alcohol and drug-related problems. Answering •yesŽ to three or four questions may indicate alcohol or drug dependence.[83]

UNCOPE

U Have you spent more time drinking or Using than you intended?

N Have you ever Neglected some of your usual responsibilities because of alcohol or drug use?

C Have you ever felt you wanted or needed to Cut down on your drinking or drug use in the past year?

O Has your family, a friend, or anyone else ever told you they Objected to your alcohol or drug use?

P Have you found yourself thinking a lot about drinking or using? (Preoccupied)

E Have you ever used alcohol or drugs to relieve Emotional discomfort, such as sadness, anger, or boredom?

Scoring: Two or more positive responses indicate possible abuse or dependence and a need for further assessment by an SUD treatment provider.[84]

does not indicate a problem, the caseworker•s best judgment always should take precedence.

What to Do When a Screen Indicates a Substance Use Disorder

If the results of a screening instrument or an in-home check indicate that a parent may have an SUD, the CPS caseworker should take the following steps:

- Ensure that the parent receives an SUD assessment from a quali“ed SUD treatment provider.
- If an SUD is present, address it in the case plans for both the parent and the child.
- Ensure that a quali“ed professional assesses the child for the impact of parental SUD or for the possibility of the child•s own use of substances.
- Coordinate service plans with the treatment professional.

Even if an SUD initially has been ruled out as an important factor in the family•s case plan, the caseworker should reassess if the family is not making progress in dealing with other issues. An unidenti“ed SUD can hamper a family•s progress for years.

Assessment

Once screening indicates that an individual may have an SUD, an assessment is the next step in a continuum of activities to address the problem. An assessment is a detailed evaluation used to determine whether treatment is needed. If so, then the assessment is

Drug Testing Parents

Physical drug testing of parents for evidence of substance use brings to the surface complicated and interrelated issues of public policy, science and technology, and ethics. Drug testing of adults has di erent goals in di erent contexts. For example, parole and probation o cers use drug testing to monitor compliance with the conditions of parole and probation; employers use them to make hiring decisions; and alcohol and drug abuse treatment programs use tests to assess whether a person is complying with the treatment plan. Parents involved with CPS who are known to have an SUD are likely to be tested as part of their alcohol and drug abuse treatment or to meet court requirements. CPS may rely on the results of drug tests to inform decisions about providing services or reuni“cation, or they may consider drug testing as a means of determining if there is an SUD. Whichever way drug testing is used, it is important for CPS caseworkers and administrators to understand the following uses and limitations:

- Drug tests do not demonstrate patterns of drug use or demonstrate if a person is abusing or is dependent on substances. Test results simply indicate the recent use of a substance and, for some substances, the amount used.
- Common drug tests do not provide accurate information about alcohol use because alcohol metabolizes quickly and is not detectable after approximately 8 hours.
- Whether drug use is detected by tests depends not only on the drug used, but also on other factors such as the characteristics of each drug, an individual•s metabolism, and the cut-o levels established by the agency requesting the test or the laboratory analyzing it.
- Drug tests are typically physically invasive procedures, which raises questions about an individual•s right to privacy.
- Individuals may be afraid to discuss problems if they believe they will be tested.
- Positive results from drug tests require that there be quali“ed and trained sta available to initiate careful and sensitive follow-up discussions with family members.

For information about how the child welfare system, SUD treatment providers, and the courts can improve screening and assessment policies and protocols, refer to *Screening and Assessment for Family Engagement, Retention, and Recovery (SAFERR)* at http://www.ncsacw.samhsa.gov/files/SAFERR.pdf.

e goal of treatment is to help individuals break the cycle of addiction and dependence so that they may learn better ways of dealing with challenges in their lives. Caseworkers should keep in mind that treatment does not equal recovery. Recovery is a lifelong process, with treatment being one of the steps. Recovery entails making lifestyle changes to regain control of one's life and accepting responsibility for one's own behavior.[90]

Research has demonstrated that SUD treatment works. A number of national studies over the past decades have shown that SUD treatment can result in abstinence from substance use, significant reduction in the abuse of substances, decreased criminal activity, and increased employment.[91] Recent studies also link SUD treatment for mothers with children in substitute care to improved child welfare outcomes, such as shorter stays in foster care for children and increased likelihood of reunification.[92] Furthermore, treatment has been shown to be cost-effective and to reduce costs in such areas as crime, health care, and unemployment.[93]

TREATMENT C

- Type and setting. An individual should be placed in the type and setting of treatment that is most appropriate for the specific problems and needs. Just as a doctor may determine that a patient should receive medication instead of surgery to correct a problem, an SUD treatment provider must make decisions about the most appropriate course of treatment for an individual. The type, length, and duration of the treatment vary depending on the type and the duration of the SUD and the individual's support system and personal characteristics. The duration of the treatment may range from weeks or months to years.

- Reassessment and modification of treatment plan. An individual's treatment and service plan should be reassessed and continually modified to ensure that the plan meets the person's evolving needs.[35]

- Involuntary treatment. An individual does not have to "hit bottom" or "want to change" in order to benefit from treatment. Involuntary or mandated treatment can be just as effective as voluntary treatment. Sanctions or enticements in the family, work, or court setting can significantly increase treatment entry, retention, and success.[36]

- Attorney involvement. In instances where the parent has an attorney, the attorney also can play a key role in the early engagement of the client in treatment. CPS caseworkers and SUD treatment providers can facilitate this by reaching out to attorneys to help them understand the treatment process and clients' needs. This helps them represent the clients better and provides a better opportunity for reunification.

- Timetables. Because of the potential conflicts between child welfare and treatment timetables, treatment should begin as soon as possible so that there is time for family reunification. Often, however, there are delays in treatment either because it is not available or the need for treatment is not determined right away. CPS caseworkers and SUD treatment providers should work together to engage clients in treatment as early as possible.

Common T

drugs, develop strategies for dealing with these situations, and build speci“c skills to address behaviors and problems that are associated with SUDs. For example, if a woman suggests that she is most likely to use cocaine after she has had a “ght with her partner, the therapist would work with her to develop more positive ways of dealing with her anger and frustration following that “ght. e treatment provider also may detail possible consequences to the individual, such as breaking parole and being forced to return to prison, as a means of changing behavior.

- Motivational enhancement treatment incorporates some elements of cognitive-behavioral treatment, but focuses on increasing and then maintaining participants• motivations for change. Rather than forcing individuals to accept that they have a problem, this approach focuses on the individual•s needs and the discrepancies between their goals and their current behaviors. is approach seeks to draw solutions from the treatment participants rather than having the solutions imposed by therapists.

- Contingency management includes both motivational enhancement treatment and an additional component of reinforcements and rewards. For example, credits may be used as a reward for established positive behaviors, such as consistent attendance in group therapy or negative urinalysis testing. ese credits then can be exchanged for items (such as baby products).

- erapeutic communityis an approach based on both cognitive-behavioral therapy and on the notion that treatment is best provided within the context of a community of individuals who have similar histories. is model was developed to provide treatment to individuals with antisocial character traits in addition to SUDs and tends to be highly confrontational. By having treatment participants confront each others• behaviors and attitudes, they learn a great deal about their own behaviors and also learn from the other participants. Often, therapeutic community models of treatment are found within the correctional system. Given its confrontational nature, a therapeutic community may not be appropriate for some individuals. For example, women who have experienced intimate partner violence likely would not react well to this treatment approach.

- Trauma-informed treatment servicesgenerally follow one or more of the above treatment theories and re”ect an understanding of trauma and its impact on SUDs and recovery.is

Model Treatment and Prevention Programs

e following Internet resources provide information about model SUD treatment and prevention interventions and their characteristics:

- e Substance Abuse and Mental Health Services Administration (SAMHSA) has compiled a list of evidenced-based programs that have prevented or reduced SUDs and other related behaviors. SAMHSA•s National Registry of Evidence-based Programs and Practices has reviewed these programs rigorously and assessed their e ectiveness. To view this list, go to http://modelprograms.samhsa.gov.
- SAMHSA•s *Guide to Evidence-Based Practices* provides listings for more than 35 websites that contain information and research on speci“c evidence-based programs and practices for the treatment or prevention of SUDs. Listings can be sorted and browsed by topic areas, target age groups, and settings. To view these listings, visit http://www.samhsa.gov/ebpWebguide/index.asp.
- e National Institute on Drug Abuse of the National Institutes of Health o ers a list of principles for substance use prevention based on a number of long-term research studies. at list can be viewed at http://www.nida.nih.gov/Infofacts/lessons.html.

approach acknowledges that a large percentage of SUD treatment participants have sustained physical, emotional, and sexual trauma in their lives and their disorder may be the result of self-medicating behaviors to deal with post-traumatic stress disorder symptoms.

- Trauma-speci“c treatment services go a step further than trauma-informed treatment services and address the impact of the speci“c trauma on the lives of participants. is approach works to facilitate trauma healing and recovery as part of the treatment services. Several integrated, trauma-speci“c, treatment models for women have been developed in recent years.[98]

- Treatment based upon the relational model of women•s development acknowledges the primacy of relationships in the lives of women and focuses upon the establishment and support of positive relationships. ese positive relationships for the treatment participant may be with the therapist or with other signi“cant “gures, especially children and spouses.

Support Services

Along with SUD treatment, supplemental services often are provided to give additional support aimed at improving treatment outcomes. e following are important support services for treatment:

- Case management services are aimed at eliminating or reducing barriers to participation in treatment and include links to housing, food, medical care, “nancial assistance, and legal services. Case management also may include problem-solving sessions to assist individuals in establishing priorities among the many demands made upon them by multiple systems.

- Twelve-step models that incorporate the 12 steps of Alcoholics Anonymous into treatment. Participants •work the stepsŽ and move through treatment by accomplishing each of the 12 steps with guidance from a sponsor and with emphasis on attendance at meetings. Spirituality or belief in a •higher powerŽ is a central component of 12> step models.

Treatment Example: Methadone Maintenance

Treatment can take many forms and can be multilayered and complex in attempting to address the nature of SUDs. For example, opioid replacement therapy is a treatment that substitutes a noneuphoria inducing and legally obtainable drug (e.g., methadone, buprenorphine) for heroin or another opiate. e treatment also provides counseling and other rehabilitation services. Methadone maintenance treatment is a type of opioid replacement therapy and is very e ective. Along with preventing illicit opiate use, methadone has been shown to be e ective in reducing criminal activity and increasing employment. Additionally, this treatment method reduces the risk of HIV-associated behaviors (e.g., needle use and sharing) and infection.[99]

Individuals engaged in methadone maintenance treatment can face heavy discrimination within the child welfare system from judges, attorneys, and caseworkers who believe the ultimate goal of treatment should be a completely drug-free individual. Stopping the methadone treatment, however, leaves the individual at a very high risk for relapse to illicit opiate use and its associated high-risk factors, including unsafe injection practices and illegal behavior in order to support a habit. All of these can signi“cantly increase the risk of abuse or neglect to children in the custody of these parents. Hence, the decision to require a medical detoxi“ from methadone must be considered carefully and based upon sound clinical principles rather than upon the stigma associated with methadone treatment.

is not uncommon. Not only do these women face the loss of a relationship, but many also face the loss of economic support. is has particular importance when the women are also mothers with young, dependent children.

Even if mothers do not have to contend with

increase safety risks. In addition, CPS caseworkers usually are required by the court to seek and to involve absent parents. In some cases, it is the fathers of the children in the child welfare system who become the focus of intervention due to the presence of mothers in treatment. In these cases, men•s roles as fathers and primary caregivers for their children warrant signi“cant attention as they struggle to provide appropriate, nurturing, and consistent parenting.

Barriers to Treatment

Most people who have SUDs do not receive treatment. According to NSDUH, approximately 23.2 million people in 2007 needed SUD treatment. Of these, 2.4 million (10.4 percent) received treatment at a specialty facility (including hospitals, drug or alcohol rehabilitation facilities, and mental health 10.193t0(ea1ent. O)6(f thes2d205d primar)-18(yencetationmains,

The Role of Child Protective Services When Substance Use Disorders Are Identified

In This Chapter

- Family assessment and case planning
- Supporting parents in treatment and recovery
- Supporting children of parents with SUDs

Once substance use disorders (SUDs) are identified as an issue to be addressed in a family•s case plan, the child protective services (CPS) caseworker needs to have a discussion with the family to understand their perceptions of the role and the impact substance abuse or dependence has in their lives. is discussion should include what can be done about the issue and how the family can be motivated to change. Since a discussion about SUDs may be met with denial and even anger toward the caseworker, a focus on the needs of the children generally will align caseworkers and parents in determining the best way to improve the situation. is chapter discusses family assessments and how they can be used in case planning, how to support parents who are in treatment and recovery, and how to assist children whose parents have SUDs.

Family Assessment and Case Planning

During the initial family assessment or investigation, the CPS caseworker identi“es the behaviors and conditions of the child, parent, and family that contribute to the risk of maltreatment, which may include a family member•s SUD. During the family assessment, the caseworker engages the family in a process designed to gain a greater understanding of family strengths, needs, and resources so that children are safe and the risk of maltreatment is reduced.[44] In particular, the caseworkers work with the family to:

- Identify family strengths that can provide a foundation for change (e.g., support systems)
- Reduce the risk of maltreatment by identifying and by addressing the factors that place children at risk
- Help the children cope with the e ects of maltreatment, parental SUDs, and other co> occurring problems.

A family-focused response to address family functioning issues is essential to an e ective case plan. Families are involved with CPS because of serious breakdowns in functioning that can be in”uenced profoundly by a family member•s SUD, as well as by the same family member•s transition to recovery. Not only must the parents• substance use be addressed, but the behavioral problems and issues that have developed for children over the span of their parents• substance use also must be resolved. To cease substance abuse and to make positive changes in their lives, it is vital for parents to move toward full acceptance of their substance abuse or addiction and its consequences. When parents address their SUDs and other issues, positive changes in family functioning can be achieved while the families also receive services through CPS.

Once the parent is in treatment, the CPS caseworker can coordinate with the SUD treatment provider to monitor progress, to develop ongoing supports, and to intervene in times of crisis. Ongoing communication allows both systems to obtain a more complete picture of the family, which will allow for the development and modi“cation of appropriate service plans.

When working with parents who are in treatment or

- € \ e relationship between the caseworker or treatment provider and the client should be collaborative with each participant bringing his own expertise.

- € \ Argument and aggressive confrontation should be avoided.[121]

e connections, realizations, new understandings, and solutions should come from the client rather than from the CPS caseworker.

For more information on motivational interviewing, go to http://www.motivationalinterview.orgor www.americanhumane.org/rmqic

SUPPORTING CHILDREN OF PARENTS WITH ; SUBSTANCE USE D

Child Protective Services and Substance Use Disorder Treatment Providers: Similarities and Differences

In This Chapter

- Areas of similarity
- Areas of difference

Just as there often is an overlap between the clients who child protective services (CPS) and substance use disorder (SUD) treatment agencies serve, there also is common ground in the structures and the principles that guide these two systems. CPS caseworkers and SUD treatment providers should understand the similarities and the differences between the two systems so that they can be

Shared Characteristics of Families Served ;

As discussed earlier, individuals with SUDs and parents who maltreat their children often have many other problems (e.g., mental illness, health issues, histories of domestic violence, poverty). ey require services that are beyond the scope of either CPS or SUD treatment agencies. Many of these problems overlap, so both CPS and SUD treatment agencies “nd themselves trying to address problems, such as a serious mental disorder, criminal records, HIV/AIDS, and limited job skills. Too often, each agency tries to tackle these varied problems on its own, overlooking opportunities to share this enormous responsibility with others.

Shared Management and Operational Challenges

CPS and SUD treatment program administrators and sta often face similar challenges in managing their agencies and operating their programs. ese challenges may be external, such as locating services that families need, coordinating with agencies that provide those services, navigating complex bureaucracies, and responding to political opinions or media coverage that portray families as unworthy of support. Other challenges are internal, such as di culties in hiring and training sta , high sta turnover and burnout, low pay, and outdated computer record-keeping systems.

To the extent that administrators and sta can design strategies that build on their common management challenges, they may ease some of these burdens. For example, both CPS and SUD treatment managers spend time locating and coordinating services, such as housing or mental health counseling, frequently for the same families. Time could be saved, and possibly outcomes improved, if managers collaborated in securing these services. In addition, managers could design joint training programs for sta from both agencies and seek continuing education units for sta who participate.

Shared Pressures to Attain Measurable Outcomes ;

Federal legislation requires both CPS and SUD treatment agencies to achieve measurable results, such as employment for adults and permanency decisions for children. erefore, managers from both systems are required to design and to monitor their programs to attain those results. is means that managers in both systems have to:

- Establish clear goals for sta
- Create internal monitoring and progress review systems
- Identify problems early and resolve them quickly.

CPS and SUD treatment program managers can share ideas for establishing processes that lead to measurable results. ey also can collaborate in designing monitoring and tracking systems in a way that provides useful information between their agencies as well as within them.

Areas of Difference

Notwithstanding these similarities, CPS and SUD treatment agencies may become confused or frustrated when trying to work together, even when they share overarching goals. e two systems di er in some fundamental ways, including how families enter programs, the choices available to families while they are participating, and the consequences for families if they cannot meet the standards required for completion. ese di erent contexts lead to di erent experiences for families involved with each system. Likewise, sta in each system face disparate experiences and challenges.

Parents can be angry or frightened when CPS

determine that their children have to be removed. When families come to the attention of CPS agencies, they often become involved with the courts, SUD treatment agencies, and other service providers. If they refuse to comply with the requirements established by these agencies, or if they cannot make adequate progress, they know they risk losing their children permanently.

In contrast, people generally enter SUD treatment voluntarily when they decide they are ready, and they leave when they want, even if they still are using substances. At times, however, courts order treatment as a condition of probation or parole. Coercive treatment has increased over the past several years, in part due to the increase in the use of drug courts, which are special courts designed for arrestees who have SUDs.

CPS and SUD treatment agencies also di er in the following ways:

- e primary focus of CPS is on the safety and well-being of children, and the primary focus of SUD treatment is on adult recovery. Sta of the two systems may see themselves as serving di erent clients, even if the clients are from the same family.
- e two systems operate under di erent laws and regulations.
- Funding for the two systems comes from di erent sources and with di erent conditions, even while often serving the same family.
- CPS caseworkers and SUD treatment providers may have di erent training, professional backgrounds and credentials, and disciplines. ey also commonly use di erent terms and have di erent de“nitions of certain terms. For example, CPS caseworkers usually do not di erentiate between substance use, abuse, or addiction. Caseworkers generally only want to know if the substance use a ects an individual•s ability to parent.
- Data collection requirements, computer systems, and management reporting requirements are often inconsistent or incompatible between the two systems.

Both systems operate within strict con dentiality guidelines and sta can be uncomfortable sharing information with each other, which can cause frustration. (See Chapter 8, *Putting It Together: Making the Systems Work for Families*, for a more detailed discussion of con“dentiality issues.)

7KLV SDJH LV LQWHQWLRQDOO\ OHIW EODQ

Online Tutorials for Knowledge-building and Cross-systems Work

e National Center on Substance Abuse and Child Welfare, an initiative of the Administration for Children and Families and the Substance Abuse and Mental Health Services Administration, has developed four free online self-tutorials to build knowledge about SUDs and child welfare and to support and facilitate cross-systems work. e tutorials are each intended for a speci“c audience: child welfare professionals, substance abuse treatment professionals, judicial o cers and attorneys in the dependency system, and legislators. A certi“cate for claiming Continuing Education Units is available upon successful completion of each tutorial. e tutorials are available at http://www.ncsacw.samhsa.gov/tutorials/index.asp.

For more information on training resources, visit http://www.childwelfare.gov/systemwide/training/.

priorities and integral elements of agency culture. If senior o cials decide to co-locate sta , they are more likely to realize that collaboration is an expected method of conducting business, not merely an agency buzzword.

Co-location can be highly e ective in helping CPS caseworkers and SUD treatment providers develop relationships that are essential to delivering comprehensive and well-organized services. It can change what are often a series of sequential referrals into concurrent discussions (case sta ngs) that bring greater expertise to case planning. Caseworker stress and burnout can be reduced if several people

4i[(CPhoncuipected)]TJ -0.0411 Tw 663 makrevaseworme tutorials are available at

CPS agencies and SUD treatment programs note that while Federal and some State laws are obstacles to information exchange, these laws did not create insurmountable barriers to collaboration.[131] This section discusses con"dentiality laws and ways to share information appropriately.

Con" dentiality Laws

Laws addressing various aspects of con"dentiality involving professional relationships, communications, and situations vary. ese laws may focus on:

- SUD treatment privacy requirements
- Mandated reporting of child abuse and neglect
- Privacy of CPS records
- Client-therapist con"dentiality statutes
- Research programs and data collection on human subjects.[132]

SUD treatment con"dentiality laws are based on the view that individuals with SUDs are more likely to seek treatment if they know that information about them will not be disclosed unnecessarily to others. Without the assurance of privacy, the fear of public disclosure of their problem possibly could prevent some individuals from obtaining needed treatment.

At times, however, there are important reasons for agencies to share information when working with the same families. Federal SUD treatment regulations specify circumstances under which it is appropriate that information be shared, including if the information relates to reports of child abuse or neglect. See Appendix I, *Con"dentiality and the Release of Substance Use Disorder Treatment Information*, for a list of circumstances in which patient record information can be released. Additionally, the Child Abuse Prevention and Treatment Act of 1974 (P.L. 93…247) requires that States allow for the public disclosure of information regarding a death, or near death, of a child when it is the result of maltreatment.

SUD treatment providers are subject to mandatory child abuse reporting laws in their States, requiring treatment sta to report incidents of suspected child abuse and neglect. However, this exemption from standard con"dentiality requirements applies only to initial reports of child abuse or neglect. It does not apply to requests or even subpoenas for additional information or records, even if the records are sought for use in civil or criminal investigations. us, patient "les and patient-identifying information protected by the Federal con"dentiality law still must be withheld from CPS agencies and the court unless there is some other authorization such as patient consent, an appropriate court order, or in some cases, a Qu ali" Service Organization Agreement (QSOA). Consent forms and QSOAs are discussed later in this chapter.

Key considerations related to the types of information that can be shared between CPS caseworkers and SUD treatment providers include:

- CPS case information Factors surrounding the case, any previous case history, the family environment, and other factors that are informative to the SUD treatment provider in conducting the assessment and in developing the treatment plan. CPS caseworkers must obtain appropriate consent to share this information.

Subpoenas

A subpoena to testify in court is not su cient to require the release of con"dential information, as speci"ed under Federal regulations related to con"dentiality, nor is a police search warrant. If subpoenaed to court to testify, an SUD treatment provider should "rst refuse, citing Federal regulations related to con"dentiality. Only with a judge•s subsequent court order that "nds a just cause to ignore this law in this particular case may a counselor testify without a client•s written consent.

- SUD screening information. Federal law and regulations allow CPS caseworkers to share with SUD treatment personnel information gathered during a screening for the purpose of referring an individual for an assessment.

- SUD diagnosis and treatment information. An SUD treatment agency may not disclose this information without written consent or court order. is is true even if the CPS agency referred the family member to the treatment program and mandated the assessment. For an example of a consent form, see Appendix J, *Sample Qualified Service Organization Agreement and Consent Form*.

- Attendance in treatment programs. SUD treatment programs may report a family member•s attendance at treatment, or their failure to attend, as long as the patient has signed a written consent that has not expired or been revoked. Attendance is often a key component of the family•s case plan.

- A treatment participant•s relapse. SUD treatment programs may report information about relapse to CPS caseworkers if that information is covered by a valid written consent signed by the patient. However, for many CPS agencies, the key information may be whether the family member is making satisfactory progress in treatment, even if relapse has occurred.

- Combined case plan. Most of the discussion between SUD treatment providers and CPS caseworkers will be permissible as long as the information discussed is covered by a valid written consent form. It is advisable to tell family members that their case will be discussed at periodic meetings or telephone calls and specify who will participate in the discussions.

If CPS caseworkers release the results of a substance abuse evaluation or any information regarding a client•s treatment, they violate Federal regulations related to con“dentiality. Everyone, not just SUD treatment providers, is bound by Federal con“dentiality statutes, and CPS caseworkers can be prosecuted for violating these laws. Caseworkers should clarify with their supervisor or their agency•s attorney any questions they may have about this statute and should document any legal advice given that pertains to this statute.

Ways to Share Information Appropriately

In order for the CPS caseworker and SUD treatment provider to communicate, it is important to obtain the client•s consent early, preferably at the time of the referral to treatment. Clients involved with CPS agencies may consent voluntarily to information disclosures in order to aid investigations of child maltreatment because their refusal to cooperate may result in losing custody of their children. However, information that has been disclosed through consent may not be used in criminal investigations or to prosecute the person. A consent form is only valid until the date, event, or condition on which it expires, or at any time when the treatment participant or client revokes consent. erefore, it is a good idea to set the expiration date far enough into the future to ensure that needed information can be retrieved by the other agency. It is permissible to have the consent form contain an end date that is "circumstances."[133] (See Appendix I, *Confidentiality and the Release of Substance Use Disorder Treatment Information*, for details about what should be included in a voluntary consent form.)

Another way that information can be shared between systems is through a QSOA. SUD treatment providers may disclose information under a QSOA without the patient•s consent. A QSOA is an agreement between two service organizations to share information about and to protect the con“dentiality of individuals they serve. A QSOA should not be confused with an MOU, which usually is an agreement between two or more organizations to provide services to a common set of clients.

A quali“ed service organization is one that provides services to the SUD treatment program. CPS agencies meet this de“nition if they provide services

Conclusion

For staff in any agency, it is easy to lose sight of the other systems and agencies that share a common client base. Families that experience SUDs and child maltreatment have needs, problems, and strengths that are diverse and complex. As a result, they often require the services of multiple agencies. It is critical that CPS caseworkers and SUD treatment providers have an understanding of the other system as well as the skills and desire to work toward a common goal. It is equally important that families are consulted in order to make certain that the collaborative structure helps them to address their SUDs and to ensure the safety and well-being of their children. With all of the parties committed to working jointly toward the same goals and being open to innovative approaches, successful outcomes can be achieved.

7KLV SDJH LV LQWHQWLRQDOO\ OHIW EOD

Endnotes

1\ U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration (SAMHSA), Oce of Applied Studies (OAS). (2009, April). Children living with substance-dependent or substance-abusing parents: 2002…2007. *e NSDUH Report* [On-line]. Available: http://oas.samhsa.gov/2k9/SAparents/SAparents.htm.

2\ DeBellis, M. D., Broussard, E. R., Herring, D. J., Wexler, S., Moritz, G., & Benitez, J. G. (2001). Psychiatric co-morbidity in caregivers and children involved in maltreatment: A pilot research study with policy implications. *Child Abuse & Neglect, 25(7)*, 923…944; Dube, S. R., Anda, R. F., Felitti, V. J., Croft, J. B., Edwards, V. J., & Giles, W. H. (2001). Growing up with parental alcohol abuse: Exposure to childhood abuse, neglect, and household dysfunction. *Child Abuse & Neglect, 25(12)*, 1627…1640; Cha n, M., Kelleher, K., & Hollenberg, J. (1996). Onset of physical abuse and neglect: Psychiatric, substance abuse, and social risk factors from prospective community data. *Child Abuse & Neglect, 20*(3), 191…203.

3 Kelleher, K., Cha n, M., Hollenberg, J., & Fischer, E. (1994). Alcohol and drug disorders among physically abusive and neglectful parents in a community-based sample. *American Journal of Public Health, 84*(10), 1586…1590.

4\ U.S. Department of Health and Human Services, Administration for Children and Families (ACF). (2008). *Child maltreatment 2006* [On-line]. Available: http://www.acf.hhs.gov/programs/cb/pubs/cm06/index.htm.

5\ U.S. Department of Health and Human Services, ACF. (1999). *Blending perspectives and building common ground. A report to Congress on substance abuse and child protection* [On-line]. Available: http://aspe.hhs.gov/hsp/subabuse99/subabuse.htm.

6\ U.S. Department of Health and Human Services, ACF. (1999); Hans, S. (1995). Diagnosis in etiologic and epidemiologic studies. In C. Jones & M. De La Rosa (Eds.), *Methodological issues: Etiology and consequences of drug abuse among women.* Washington, DC: U.S. Department of Health and Human Services, National Institutes of Health (NIH), National Institute on Drug Abuse (NIDA); Tarter, R., Blackson, T., Martin, C., Loeber, R., & Moss, H. (1993). Characteristics and correlates of child discipline practices in substance abuse and normal families. *American Journal on Addictions, 2*(1), 18…25; Kumpfer, K. L., & Bayes, J. (1995). Child abuse and drugs. In J. H. Jaæ (Ed.), *e encyclopedia of drugs and alcohol* (Vol. 1, pp. 217…222). New York, NY: Simon & Schuster.

7 U.S. Department of Health and Human Services, ACF. (1999).

8\ Young, N. K., & Gardner, S. L. (2002). Navigating the pathways: Lessons and promising practices in linking alcohol and drug services with child welfare. *Technical Assistance Publication (TAP) 27*. Rockville, MD: U.S. Department of Health and Human Services, SAMHSA.

9\ American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders, 4th Ed., Text Revision (DSM-IV-TR)*. Washington, DC: Author.

10\ U.S. Department of Health and Human Services, SAMHSA. (2008). *Results from the 2007 National Survey on Drug Use and Health: National findings* [On-line]. Available: http://www.oas.samhsa.gov/nsduh/2k7nsduh/2k7Results.cfm#TOC.

domestic violence [On-line]. Available: http://www.childwelfare.gov/pubs/usermanuals/domesticviolence/domesticviolence.pdf

29 \ Cunradi, C. B., Caetano, R., & Schafer, J. (2002). Alcohol-related problems, drug use, and male partner violence severity. *Alcoholism: Clinical and Experimental Research, 26*(4), 493…500; Brickley, M. R., & Shephard, J. P. (1995).e relationship between alcohol intoxication, injury severity and Glasgow coma scores in assault patients. *Injury, 26*(5), 311…314; Shephard, J. P., & Brickley, M. R. (1996). e relationship between alcohol intoxication, stressors and injury in urban violence. *British Journal of Criminology, 36*(4), 546…566.

30 \ Najavits, L. M. (2001). *Seeking safety: A treatment manual for PTSD and substance abuse.* New York, NY: Guilford Press.

31 \ Kessler, R. C., Sonnega, A., Bromet, E., Hughes, M., & Nelson, C. B. (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry*, *52*(12), 1048…1060.

32 \ Resnick, H. S., Kilpatrick, D. G., Dansky, B. S., Saunders, B. E., & Best, C. L. (1993). Prevalence of civilian trauma and posttraumatic stress disorder in a representative sample of women. *Journal of Consulting and Clinical Psychology*, *61*(6), 984…991.

33 \ Smyth, N. (1998). Exploring the nature of the relationship between poverty and substance abuse: Knowns and unknowns.

html#3.

45\\ O ce of National Drug Control Policy. (2006). *Drug Endangered Children (DEC)* [On-line]. Available: http://www.whitehousedrugpolicy.gov/enforce/dr_endangered_child.html.

46 \ Otero, C., et al. (2004).

47 \ National Association of Counties. (2005). *e meth epidemic in America: Two surveys of U.S. counties: e criminal e ect of meth on communities & the impact of meth on children* [On-line]. Available: http://www.naco.org.

48 \ U.S. Department of Health and Human Services, SAMHSA, OAS. (2009).

49 \ U.S. Department of Health and Human Services, ACF. (1999).

50 \ Zuckerman, B. (1994). E ects on parents and children. In D. Besharov (Ed.), *When drug addicts have children.* Washington, DC: Child Welfare League of America.

51 \ Child Abuse Prevention and Treatment Act, 42 U.S.C. 5106g, §Sec.111-2.

52 Child Abuse Prevention and Treatment Act, 42 U.S.C. 5106g, §Sec.111-2.

53 \ U.S. Department of Health and Human Services, SAMHSA. (2008).

54 \ Landdeck-Sisco, J. (1997). *Children with prenatal drug and/or alcohol exposure* [On-line]. Available: http://www.archrespite.org/archfs49.htm.

55 \ U.S. Department of Health and Human Services, SAMHSA. (2007). *Treatment Episode Data Set (TEDS) 1995–2005: National admissions to substance abuse treatment services* [On-line]. Available: http://wwwdasis.samhsa.gov/teds05/tedsad2k5web.pdf.

56 \ U.S. Department of Health and Human Services, SAMHSA. (2002). *Pregnant women in substance abuse treatment. e DASIS report* [On-line]. Available: http://www.oas.samhsa.gov/2k2/pregTX/pregTX.htm.

57 \ Child Welfare Information Gateway. (2006). *Prenatal drug use as child abuse* [On-line]. Available: www.childwelfare.gov/systemwide/laws_policies/statutes/drugexposedall.pdf

58 \ University of California-Santa Barbara, School of Sociology. (n.d.). *Ects of drugs taken during pregnancy* [On-line]. Available: http://www.soc.ucsb.edu/sexinfo/?article=pregnancy&id=603.

59 \ National Organization on Fetal Alcohol Syndrome. (2004). *Frequently asked questions* [On-line]. Available: http://www.nofas.org/faqs.aspx?id=9.

60 \ McCreight, B. (1997). *Recognizing and managing children with fetal alcohol syndrome/fetal alcohol e ects: A guidebook* (pp. 9…15). Washington, DC: Child Welfare League of America.

61 \ Phibbs, C. S., Bateman, D. A., & Schwartz, R. M. (1991). e neonatal costs of maternal cocaine use. *Journal of the American Medical Association, 266*(11), 1521…1526; Zuckerman, B., Frank, D. A., Hingson, R., Amaro, H., & Levenson, S. M. (1989). Ects of maternal marijuana and cocaine use on fetal growth. *New England Journal of Medicine, 320*(14), 762…768; Canadian Health Network, Canadian Centre on Substance Abuse. (2004). *What are the ects of alcohol and other drugs during pregnancy?* [On-line]. Available: http://www.canadian-health-network.ca.

62 \ Noland, J. S., Singer, L. T., & Arendt, R. E. (2003). Executive functioning in preschool-aged children prenatally exposed to alcohol, cocaine, and marijuana. *Alcoholism: Clinical & Experimental Research, 27*(4), 647…656.

63 \ Vogel, A. (1997). Neurodevelopment: Cocaine wreaks subtle damage on developing brains. *Science, 278*(5335), 38…39.

64 \ Otero, C., et al. (2004).

65 \ Irwin, K. (1995). Ideology, pregnancy, and drugs: Di erences between crack-cocaine, heroin, and methamphetamine users. *Contemporary Drug Problems, 22*(4), 613-638.

66 \ Join Together Online. (2005). *Meth science not stigma: Open letter to the media* [On-line]. Available: http://www.jointogether.org.

67 \ Finfgeld, D. (2001). Emergent drug abuse resolution models and their implications for childbearing and childrearing women. *Health Care for Women International, 22*(8), 723…733.

68 \ U.S. Department of Health and Human Services, ACF. (1999).

69 \ Goldman, J., & Salus, M. K. (2003). *A coordinated response to child abuse and neglect: e foundation for practice* [On-line]. Available: http://www.childwelfare.gov/pubs/usermanuals/foundation/index.cfm; Sullivan, S. (2000). *Child neglect: Current de nitions and models—A review of child neglect research, 1993–1998*. Ottawa, Canada: National Clearinghouse on Family Violence; Perry, B. D. (1997). *Incubated*

in terror: Neurodevelopmental factors in the 'cycle of violence' [On-line]. Available: http://www.childtrauma.org/CTAMATERIALS/incubated.asp; Kraemer, G. W. (1992). A psychobiological theory of attachment. *Behavioral and Brain Sciences, 15*(3), 493…511.

70\\ Shonko , J., & Phillips, D. (Eds.). (2000). *From neurons to neighborhoods: e science of early childhood development.* Washington, DC: National Academy Press; Kagan, J. (1999). e role of parents in children•s psychological development. *Pediatrics, 104*(1), 164…167.

71 \ Boszormenyi-Nagy, I., & Spark, G. M. (1973). *Invisible loyalties: Reciprocity in intergenerational family therapy.* New York, NY: Harper & Row.

72 \ Kilpatrick, D., Acierno, R., Saunders, B., Resnick, H., & Best, C. (2000). Risk factors for adolescent substance abuse and dependence: Data from a national sample. *Journal of Consulting and Clinical Psychology, 68*(1), 19…30.

73 \ National Research Council, Panel on Research on Child Abuse and Neglect. (1993). *Understanding child abuse and neglect*. Washington, DC: National Academy Press; Dembo, R., Dertke, M., La Voie, L., Borders, S., Washburn, M., & Schmeidler, J. (1987). Physical abuse, sexual victimization and illicit drug use: A structural analysis among high-risk adolescents. *Journal of Adolescence, 10*(1), 13…34.

74 \ National Research Council, Panel on Research on Child Abuse and Neglect. (1993).

75 \ McCauley, J., Kern, D. E., Kolodner, K., Dill, L., Schroeder, A. F., DeChant, H. K., et al. (1997).

89 \ Feild, T., & Wintefeld, A. P. (2003). *Tough problems, tough choices: Guidelines for needs-based service planning in child welfare* [On-line]. Available: http://www.americanhumane.org/protecting-children/resources/casey-decision-making-guidelines/products.html.

90 \ Breshears, E. M., et al. (2004).

91 \ National Association of State Alcohol and Drug Abuse Directors. (2001, March). *Alcohol and other drug treatment e ectiveness: A review of state outcome studies* [On-line]. Available: http://www.nasadad.org/resource.php?base_id=91; Hubbard, R. L., Craddock, S. G., Flynn, P. M., Anderson, J., & Etheridge, R. M. (1997). Overview of one-year follow-up outcomes in DATOS. *Psychology of Addictive Behavior*, *11*(4), 261…278; University of Chicago, National Opinion Research Center. (1997). *National treatment improvement evaluation study. Final report* [On-line]. Available: http://www.icpsr.umich.edu/SAMHDA/NTIES/NTIES-PDF/ntiesfnl.pdf; Gerstein, D. R., Johnson, R. A., Harwood, H., Fountain, D., Suter, N., & Malloy, K. (1994). *Evaluating recovery services: e California Drug and Alcohol Treatment Assessment (CALDATA)*. Sacramento, CA: California Department of Alcohol and Drug Programs; Hubbard, R., Mardsen, M., Rachal, J., Harwood, H., Cavanaugh, E., & Ginzburg, H. (1989). *Drug abuse treatment: A national study of e ectiveness*. Chapel Hill, NC: University of North Carolina Press; Simpson, D. (1981). Treatment for drug abuse: Follow-up outcomes and length of time spent. *Archives of General Psychiatry, 38*(8), 875…880.

92 \ Green, B. G., Rockhill, A. M., & Furrer, C. J. (2007). Does substance abuse treatment make a di erence for child welfare case outcomes? A statewide longitudinal analysis. *Children and Youth Services Review, 29*(4), 460…473.

93 \ Gerstein, D. R., et al. (1994).

94 \ U.S. Department of Health and Human Services, NIH, NIDA. (2005b).

95 \ U.S. Department of Health and Human Services, NIH, NIDA. (2005b).

96 \ U.S. Department of Labor. (2006). *Treatment* [On> line]. Available: http://www.dol.gov/asp/programs/drugs/workingpartners/sab/treatment.asp.

97 \ Breshears, E. M., et al. (2004).
NIH, NIDA. (2005b).

line]. Available: http://oas.samhsa.gov/2k3/dualTX/dualTX.htm.

110 \ U.S. Department of Health and Human Services, NIH, NIDA. (2005c). *NIDA InfoFacts: Treatment methods for women* [On-line]. Available: http://www.drugabuse.gov/Infofacts/TreatWomen.html.

117 \

111 \ U.S. Department of Health and Human Services, SAMHSA. (2008).

112 \ U.S. Department of Health and Human Services, SAMHSA, OAS. (2007). *e NSDUH Report: Substance use treatment among women of childrearing age* [On-line]. Available: http://www.oas.samhsa.gov/2k7/womenTX/womenTX.cfm.

113 \ U.S. Department of Health and Human Services, SAMHSA. (2008).

114 \ DePan“lis, D., & Salus, M. K. . *Child protective services: A guide for caseworkers* [On-line]. Available: http://www.childwelfare.gov/pubs/usermanuals/cps/cps.pdf.

115 \ National Family Preservation Network. (2005). *North Carolina Family Assessment Scale* [On-line]. Available: http://www.nfpn.org/preservation/assessment_tool.php.

116 \ Breshears, E. M., et al. (2004).

117 \ North Carolina Division of Social Services, & Family and Children•s Resource Program. (2005). *Crafting a table: Input for the Program. (2005).*

[134] U.S. Department of Health and Human Services, SAMHSA. (1999c). *Welfare reform and substance abuse treatment confidentiality: General guidance for reconciling need to know and privacy*. Rockville, MD: Author.

[135] U.S. Department of Health and Human Services, ACF. (2005). *Information memorandum ACYF-CB-IM-05-07* [On-line]. Available: http://www.acf.hhs.gov/programs/cb/laws_policies/policy/im/im0507.htm.

needs that must be addressed, and the strengths on which to build.

Family Group Conferencing … a family meeting model used by CPS agencies to optimize family strengths in the planning process. is model brings the family, extended family, and others important in the family•s life (e.g., friends, clergy, neighbors) together to make decisions regarding how best to ensure the safety of the family members.

Family Unity Model … a family meeting model used by CPS agencies to optimize family strengths in the planning process. is model is similar to the Family Group Conferencing model.

Full Disclosure … CPS information to the family regarding the steps in the intervention process, the requirements of CPS, the expectations for the family, the consequences if the family does not ful" the expectations, and the rights of the parents to ensure that the family completely understands the process.

Guardian ad Litem … a lawyer or lay person who represents a child in juvenile or family court. Usually this person considers the best interest of the child and may perform a variety of roles, including those of independent investigator, advocate, advisor, and guardian for the child. A lay person who serves in this role is sometimes known as a court-appointed special advocate or CASA.

Habituation … the result of repeated consumption of a drug that produces psychological, but not physical, dependence. e psychological dependence produces a desire (not a compulsion) to continue taking drugs for the sense of improved well-being.

Home Visitation Programs … prevention programs that o er a variety of family-focused services to pregnant women and families with new babies. Activities frequently encompass structured visits to the family•s home and may address positive parenting practices, nonviolent discipline techniques, child development, maternal and child health, available services, and advocacy.

Immunity …established in all child abuse laws to protect reporters from civil law suits and criminal prosecution resulting from “ling a report of child abuse and neglect.

Initial Assessment or Investigation … the stage of the CPS case process during which the CPS caseworker determines the validity of the child maltreatment report, assesses the risk of maltreatment, determines if the child is safe, develops a safety plan if needed to ensure the child•s protection, and determines services needed.

Intake … the stage of the CPS case process in which the CPS caseworker screens and accepts reports of child maltreatment.

Interview Protocol … a structured format to ensure that all family members are seen in a planned strategy, that community providers collaborate, and that information gathering is thorough.

Involuntary Commitment … process by which patients who have not committed any crime are brought to SUD treatment against their wishes by relatives, police, or through a court proceeding. Also known as •protective custodyŽ or •emergency commitment.Ž

Juvenile and Family Courts… established in most States to resolve con”ict and to otherwise intervene ht to SUD 8 Tw T*ectsu0 -ectment.

Service Provision … the stage of the CPS casework process during which CPS and other service providers o er speci“c services to reduce the risk of maltreatment.

Sexual Abuse … inappropriate adolescent or adult sexual behavior with a child. It includes fondling a child•s genitals, making the child fondle the adult•s genitals, intercourse, incest, rape, sodomy, exhibitionism, sexual exploitation, or exposure to pornography. To be considered child abuse, these acts have to be committed by a person responsible for the care of a child (for example a babysitter, a parent, or a day care provider) or related to the child. If a stranger commits these acts, it would be considered sexual assault and handled solely by the police and criminal courts.

Substance Abuse … a pattern of substance use resulting in clinically signi“cant physical, mental, emotional, or social impairment or distress, such as failure to ful“ll major role responsibilities, or use in spite of physical hazards, legal problems, or interpersonal and social problems.

Substance Dependence … see •addiction.Ž

Substance Use … consumption of low or infrequent doses of alcohol and other drugs, sometimes called experimental, casual, or social use, such that damaging consequences may be rare or minor.

Substance Use Disorder (SUD) … a medical condition that includes the abuse of or addiction to (or dependence on) alcohol or drugs.

Substantiated … an investigation disposition concluding that the allegation of maltreatment or risk of maltreatment was supported or founded by State law or State policy. A CPS determination means that credible evidence exists that child abuse or neglect has occurred.

System of Care … a comprehensive continuum of child welfare, SUD, and other support services coordinated to meet the multiple, evolving needs of clients.

Tertiary Prevention … treatment e orts geared to address situations in which child maltreatment already has occurred, with the goals of preventing child maltreatment from occurring in the future and of avoiding the harmful e ects of child maltreatment.

Tolerance … a state in which the body•s tissue cells adjust to the presence of a drug in given amounts and eventually fail to respond to ordinarily e ective dosages. Consequently, increasingly larger doses are necessary to produce desired e ects.

Treatment … the stage of the child protection case process during which speci“c services are delivered by CPS and other providers to reduce the risk of maltreatment, support families in meeting case goals, and address the e ects of maltreatment.

Universal Prevention … activities and services directed toward the general public with the goal of stopping maltreatment before it starts. Also referred to as •primary prevention.Ž

Unsubstantiated (also Not Substantiated) … an investigation disposition that determines that there is not su cient evidence under State law or policy to conclude that the child has been maltreated or is at risk of maltreatment. A CPS determination means that credible evidence does not exist that child abuse or neglect has occurred.

Withdrawal … symptoms that appear during the process of stopping the use of a drug that has been taken regularly.

Resource Listings of Selected National Organizations Concerned with Child Maltreatment and/or Substance Use Disorders

e following are several representatives of the many national organizations and groups dealing with various aspects of child maltreatment and substance use disorders. Visit http://www.childwelfare.gov view a more comprehensive list of resources and visit http://www.childwelfare.gov/organizations/index.cfm to search an organization database. Inclusion on this list is for information purposes only and does not constitute an endorsement by the Co on Child Abuse and Neglect or the Children•s Bureau.

National And Federal Substance Use ; Disorder Organizations ;

Addiction Technology Transfer Centers ;

address: National Oce
University of Missouri…Kansas City
5100 Rockhill Road
Kansas City, MO 64110-2499

phone: (816) 235-6888

fax: (816) 235-6580

email: no@nattc.org

website: http://www.nattc.org

A nationwide, multidisciplinary resource that draws upon the knowledge, experience, and latest work of recognized experts in the “eld of addictions.

Community Anti-Drug Coalitions of America ;

address: 625 Slaters Lane, Suite 300
Alexandria, VA 22314

phone: (800) 54-CADCA

fax: (703) 706-0565

email: info@cadca.org

website: http://cadca.org

Builds and strengthens the capacity of community coalitions to create safe, healthy, and drug-free communities. Supports its members with technical assistance and training, public policy, media strategies and marketing programs, conferences, and special events.

Join Together Online

address: 715 Albany Street, 580-3rd Floor
Boston, MA 02118

phone: (617) 437-1500

fax: (617) 437-9394

email: info@jointogether.org

website: http://www.jointogether.org

Supports community-based e orts across the country to reduce, prevent, and treat substance use disorders. Focuses attention on strengthening community capacity to expand the demand for and supply of high-quality substance use disorder treatment.

Substance Abuse and Mental Health Services Administration

address: 1 Choke Cherry Road
Rockville, MD 20857

phone: (240) 276-2000

fax: (240) 276-2010

website: http://www.samhsa.gov

Seeks to improve the quality and availability of prevention, treatment, and rehabilitative services in order to reduce illness, death, disability, and cost to society resulting from substance use disorders and mental illnesses.

Child Welfare Organizations

American Humane Association
Children•s Division

address: 63 Inverness Drive, East
Englewood, CO 80112-5117

phone: (303) 792-9900

fax: (303) 792-5333

website: www.americanhumane.org

Conducts research, analysis, and training to help public and private agencies respond to child maltreatment.

American Professional Society on the Abuse of Children

address: 350 Poplar Avenue
Elmhurst, IL 60126

phone: (630) 941-1235
(877) 402-7722

fax: (630) 359-4274

email: apsac@apsac.org

website: www.apsac.org

Provides professional education, promotes research to inform e ective practice, and addresses public policy issues. Professional membership organization.

American Public Human Services Association

address: 810 First Street, NE, Suite 500
Washington, DC 20002-4267

phone: (202) 682-0100

fax: (202) 289-6555

website: www.aphsa.org

Addresses program and policy issues related to the administration and delivery of publicly funded human services. Professional membership organization.

AVANCE Family Support and Education Program

address: 118 N. Medina
San Antonio, TX 78207

phone: (210) 270-4630

fax: (210) 270-4612

website: www.avance.org

Operates a national training center to share and disseminate information, materials, and curricula to service providers and policymakers interested in supporting high-risk Hispanic families.

Child Welfare League of America

address:

National Black Child Development Institute ;

address: 1313 L Street, NW
Suite 110
Washington, DC 20005-4110

phone: (202) 833-2220

fax: (202) 833-8222

email: moreinfo@nbcdi.org

website: www.nbcdi.org

Operates programs and sponsors a national training conference through Howard University to improve and protect the well-being of African-American children.

National Children•s Advocacy Center

address: 210 Pratt Avenue
Huntsville, AL 35801

phone: (256) 533-KIDS

fax: (256) 534-6883

website: http://www.nationalcac.org

Provides prevention, intervention, and treatment services to physically and sexually abused children and their families within a child-focused team approach.

National Indian Child Welfare Association

address: 5100 SW Macadam Avenue, Suite 300
Portland, OR 97239

phone: (503) 222-4044

fax: (503) 222-4007

website: www.nicwa.org

Disseminates information and provides technical assistance on Indian child welfare issues. Supports community development and advocacy to facilitate Tribal responses to the needs of families and children.

National Resource Center for Child Protective Services

address: 925 #4 Sixth Street, NW
Albuquerque, NM 87102

phone: (505) 345-2444

fax: (505) 345-2626

website: http://www.nrccps.org

Focuses on building State, local, and Tribal capacity through training and technical assistance in child protective services, including meeting Federal requirements, strengthening programs, eligibility for the Child Abuse Prevention and Treatment Act grant, support to State Liaison Officers, and collaboration with other national resource centers.

CHILD ABUSE PREVENTION ORGANIZATIONS

National Alliance of Children•s Trust and Prevention Funds

address:

Prevent Child Abuse America ;

address:

State Telephone Numbers for Reporting Child Maltreatment

Each State designates speci“c agencies to receive and investigate reports of suspected child abuse and neglect. Typically, this responsibility is carried out by child protective services (CPS) within a Department of Social Services, Department of Human Resources, or Division of Family and Children Services. In some States, police departments also may receive reports of child abuse or neglect.

Many States have local or toll-free telephone numbers for reporting suspected maltreatment. The reporting party must be calling from the same State where the child is allegedly being maltreated for most of the following numbers to be valid.

For States not listed, or when the reporting party resides in a di erent State from the child, please call Childhelp, 800-4-A-Child (800-422-4453), or your local CPS agency. States may occasionally change the telephone numbers listed below. To view the most current contact information, including State Web Addresses, visit http://www.childwelfare.gov/pubs/reslist/rl_dsp.cfm?rs_id=5&rate_chno=11-11172

Alabama (AL)
334-242-9500

Alaska (AK)
800-478-4444

Arizona (AZ)
888-SOS-CHILD
(888-767-2445)

Arkansas (AR)
800-482-5964

Colorado (CO)
303-866-5932

Connecticut (CT)
800-842-2288

Delaware (DE)
800-292-9582

District of Columbia (DC)
202-671-SAFE (7233)

Florida (FL)
800-96-ABUSE
(800-962-2873)

Hawaii (HI)
808-832-5300

Idaho (ID)
800-926-2588

Illinois (IL)
800-252-2873

Indiana (IN)
800-800-5556

Iowa (IA)
800-362-2178

Kansas (KS)
800-922-5330

Kentucky (KY)
800-752-6200

Maine (ME)
800-452-1999
800-963-9490 (TTY)

Massachusetts (MA)

Mississippi (MS)
800-222-8000
601-359-4991

Missouri (MO)
800-392-3738
573-751-3448

Montana (MT)
866-820-KIDS (5437)

Nebraska (NE)
800-652-1999

Nevada (NV)
800-992-5757

New Hampshire (NH)
800-894-5533
603-271-6556

New Jersey (NJ)
877-652-2873
800-835-5510 (TDD/TTY)

New Mexico (NM)
800-797-3260
505-841-6100

New York (NY)
800-342-3720
518-474-8740
800-369-2437 (TDD)

Oklahoma (OK)
800-522-3511

Pennsylvania (PA)
800-932-0313

Puerto Rico (PR)
800-981-8333
787-749-1333

Rhode Island (RI)
800-RI-CHILD
(800-742-4453)

Diagnostic and Statistical Manual of Mental Disorders Criteria [1]

Criteria for Substance Dependence

A maladaptive pattern of substance use, leading to clinically signi“cant impairment or distress, as manifested by three (or more) of the following, occurring at any time in the same 12-month period:

1. \ Tolerance, as de“ned by either of the following:
    a. \ A need for markedly increased amounts of the substance to achieve intoxication or desired e ect
    b.\ \Markedly diminished e ect with continued use of the same amount of the substance.
2. \ Withdrawal, as manifested by either of the following:
    a. \ e characteristic withdrawal syndrome for the substance (refer to criteria A and B of the criteria sets for withdrawal from the speci“c substances).
    b.\ \ e same (or a closely related) substance is taken to relieve or avoid withdrawal symptoms.
3. \ e substance often is taken in larger amounts or over a longer period than was intended.
4. \ ere is a persistent desire or unsuccessful e orts to cut down or control substance use.
5. \ A great deal of time is spent in activities necessary to obtain the substance (e.g., visiting multiple doctors or driving long distances), use the substance (e.g., chain-smoking), or recover from its e ects.
6. \ Important social, occupational, or recreational activities are given up or reduced because of substance use.
7. \ e substance use is continued despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by the substance (e.g., current cocaine use despite recognition of cocaine-induced depression, or continued drinking despite recognition that an ulcer was made worse by alcohol consumption).

[1] American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders (4th ed.) Text Revision (DSM-IV-TR).* Washington, DC: Author.

Specify if:

a. \ With physiological dependence„evidence of tolerance or withdrawal (i.e., either Item 1 or 2 is present).

b. Without physiological dependence„no evidence of tolerance or withdrawal (i.e., neither Item 1 nor 2 is present).

Criteria for Substance Abuse

A maladaptive pattern of substance use leading to clinically signi“cant impairment or distress, as manifested by one (or more) of the following, occurring within a 12-month period:

1. \ Recurrent substance use resulting in a failure to ful“ll major role obligations at work, school, or home (e.g., repeated absences or poor work performance related to substance use; substance-related absences, suspensions, or expulsions from school; neglect of children or household).

2. \ Recurrent substance use in situations in which it is physically hazardous (e.g., driving an automobile or operating a machine when impaired by substance use).

3. \ Recurrent substance-related legal problems (e.g., arrests for substance-related disorderly conduct).

4. \ Continued substance use despite having persistent or recurrent social or interpersonal problems caused or exacerbated by the e ects of the substance (e.g., arguments with spouse about consequences of intoxication, physical ghts).

e symptoms never have met the criteria for substance dependence for this class of substance.

Commonly Abused Substances

is appendix presents information about some of the most commonly abused types of substances and the drugs a liated with those types. A table provides additional information on commonly abused substances, including street names and methods of use.

Types of Commonly Abused Substances

Cannabinoids (e.g., marijuana, hashish) can produce feelings of euphoria, anxiety, or depression as well as distort perception and slow reaction time. Marijuana, the most commonly used illicit drug, has been associated with automobile and industrial accidents as well as physical ailments, most notably cancer[1]. It is usually smoked, making its health risks similar to those of tobacco in terms of pulmonary and cardiac e ects.

Club Drugs (e.g., GHB, ketamine, MDMA, " unitrazepan, yaba) include a variety of drugs from other drug categories (e.g., hallucinogens, depressants). e name is derived from the fact these substances often are used by younger people at nightclubs and parties. Some club drugs, gamma hydroxyl butyrate (GHB) and Rohypnol in particular, have gained notoriety for use in drug-assisted sexual assault cases and, therefore, are referred to as "predatory drugs." Certain club drugs, such as ketamine, have medical or veterinary uses, but are used in a significantly di erent quantity or by a population other than for whom they are intended. Yaba, which means "crazy medicine" in Thai, is a combination of methamphetamine and ca eine and is becoming increasingly available at rave parties. Like methamphetamine, use of yaba can result in a rapid heart rate and damage to the small blood vessels in the brain, which can lead to stroke. Its use also can lead to violent behavior, paranoia, confusion, or insomnia.

Depressants (e.g., barbiturates) include some drugs that are prescribed to reduce anxiety or act as a sedative or anticonvulsant. Depressants are used illicitly to produce feelings of well-being and to lower inhibitions. Signs of use include fatigue, confusion, and impaired coordination and memory. Alcohol, the most commonly abused substance, is categorized as a depressant.

Hallucinogens (e.g., acid, mescaline, psilocybin, phencyclidine) have no known medical use and are illegal. ese substances produce altered states of perception and feeling. Users often are disoriented or inattentive. e e ects of hallucinogens are unpredictable and depend on several factors, including the user's personality, the surroundings in which they are used, the quantity taken, and the drug's purity.

[1] U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration. (2003). *Marijuana: Facts for teens* [On-line]. Available: http://www.nida.nih.gov/MarijBroch/Marijteens.html.

Users may experience a •bad tripŽ that can include terrifying thoughts and feelings, fear of insanity or death, and deep despair. An observable, long-term e ect for some users is persistent perception disorder, which is commonly referred to as "flashbacks.

Inhalants (e.g., aerosol sprays, nitrous oxide, butyl nitrate) usually are legal and readily available household and commercial products whose chemical vapors are inhaled to produce mind-altering e ects. Observable e ects of use include runny nose, watery eyes, and headaches. Users can ingest the substances by inhaling directly from product containers, sni ng a cloth saturated with the substance, or sni ng the substance from a plastic bag that is placed over the nose and mouth. Deeply inhaling vapors or using large amounts over a short time may result in disorientation, violent behavior, unconsciousness, or even death. High concentrations of inhalants can cause su ocation by displacing oxygen in the lungs. One of the signi“cant factors in the use of inhalants is their accessibility, particularly for children. National surveys indicate inhaling dangerous products , is becoming a widespread problem.

Opioids and Narcotic Pain Relievers (e.g., heroin, morphine, oxycodone, hydrocodone) are used illegally for their euphoric e ects. Many opioids and narcotic pain relievers originally were developed to relieve pain, and doctors still prescribe some for that purpose. e pain of withdrawal from heroin and other opioids is made worse by the fact that these drugs deaden pain. erefore, individuals in withdrawal may experience pain they did not feel while using opiates and may not be able to deal with pain as they normally would. Signs of use include needle marks as well as decreased pulse and respiration rates.

Stimulants (e.g., cocaine, amphetamines, methamphetamine) can produce e ects such as increased alertness, over-activity, depression, and insomnia. Cocaine is a powerful stimulant that primarily a ects the dopamine system, the part of the brain that regulates feelings of pleasure and excitement. Cocaine use can cause violent or hypersexual behavior, paranoid thinking, and agitation or anxiety. Prenatal exposure to cocaine can cause premature labor, low birth weight, and feosurh.035o9xy
ıal exposure to cocain ploidd to problugs

ıng er* [(lafabeir 0(ech ten, a* [((citemes.)]TJ EMC /P1 <<

Selected Substance Use Disorder Screening Instruments			
Instrument	Purpose	Summary	For More Information
Alcohol Use Disorders Identi“cation Test (AUDIT) [1]	Designed to identify individuals whose alcohol use has become a danger to their health. Includes three subscales that assess amount and frequency of drinking, alcohol dependence, and problems caused by alcohol.	€ Ten questions that can be self-administered or asked by another person € Takes about 1 minute to complete. € Targeted at adults. € Free except for training materials.	omas Babor Alcohol Research Center University of Connecticut 263 Farmington Avenue Farmington, CT 06030-2103
CAGE Questionnaire [2]	Related to drinking behavior.	€ Four questions that can be self-administered or asked by another person € Targeted at individuals 16 years of age or older. € Questions can be incorporated into other questionnaires.	Available through numerous publications, websites, and treatment and prevention programs, including: http://pubs.niaaa.nih.gov/publications/Assesing%20

Selected Substance Use Disorder Screening Instruments			
Instrument	Purpose	Summary	For More Information
Michigan Alcoholism Screening Test (MAST)	Designed to screen for lifetime alcoholism-related problems.	€ Twenty-“ve questions that can be self-administered or asked by another person. Shorter version also available. € Takes 5 minutes to administer. € Targeted at adults. € Minor cost for original,	Melvin L. Selzer, M.D.

Triage Assessment for Addictive Disorders (TAAD)	Designed for both drug and alcohol use in face-to-face interviews where time commitment is minimal.	€ irty questions. € Takes 12…13 minutes to administer and score. € Can be administered by anyone with good interviewing skills. € Requires expertise to score. € Must be purchased.	Norman G. Ho mann, Ph.D. Evince Clinical Assessments P.O. Box 17305 Smith“eld, RI 02917 (800) 755-6299 http://www.evinceassessment.com
TWEAK Alcohol Screen	Developed and validated for women. Recommended by the California Institute of Mental Health.	€ € € Evince C0.12olnsann, Ph.D. € €	

9. Yes R No R The client has used the following in the last 12 months:

Marijuana/Hashish R

Memorandums of Understanding [1]

What is a Memorandum of Understanding (MOU)?

An MOU is a written agreement that serves to clarify relationships and responsibilities between two or more organizations that share services, clients, and resources.

Why is it important to have an MOU?

MOUs help strengthen community partnerships by delineating clear roles between individuals, agencies, and other groups. Communities with MOUs report that the strengthened partnerships resulted in enhanced services for children and families.

What is actually included in an MOU?

Generally, MOUs can address a variety of issues and topics. Content areas to consider including in an MOU are:

- Clari“cation of agency roles;
- Referrals across agencies;
- Assessment protocols;
- Parameters of con dentiality;
- Case management intervention;
- Interagency training of sta ;
- Agency liaison and coordination;
- Process for resolving interagency con icts;
- Periodic review of the MOU.

[1] Bragg, H. L. (2003). *Child protection in families experiencing domestic violence* [On-line]. Available: http://www.childwelfare.gov/pubs/usermanuals/domesticviolence/.

How do we know our community is ready to develop an MOU?

Communities that are concerned about reducing the incidence of child maltreatment are excellent candidates for creating an MOU. In communities that are experiencing strained relationships between potential partners, the process of writing an MOU provides a unique opportunity to address misperceptions and di erences and to work jointly to resolve gaps in service delivery.

What strategies should we undertake as we begin the MOU process?

Depending on existing relationships within communities, one strategy may include inviting key supporters to

Confidentiality and the Release of Substance Use Disorder Treatment Information

e Comprehensive Alcohol Abuse and Alcoholism Prevention, Treatment, and Rehabilitation Act (1970) and the Drug Abuse O ce and Treatment Act (1972) regulate the disclosure and confidentiality of information by substance use disorder treatment programs that receive Federal assistance. Generally, a provider cannot release any information that identi“es an individual in the program and cannot acknowledge the presence of an individual in the treatment program. e following are exceptions under which client information can be released:

- € \ It will be used in internal communications between or among those with a legitimate interest who need the information in connection with their duties that arise out of the provision of diagnosis, treatment, or referral for treatment of substance use disorders if the communications are within the program or between a program and an entity that has direct administrative control over the program.
- € \

… A statement regarding revocation of consent;

… e date, event, or condition upon which the consent will expire;

… e signature of the patient;

… e date on which the consent is signed.[1]

Adapted from U.S. Department of Justice, Office of Juvenile Justice and Delinquency Prevention. (2000). Legal issues. Juvenile Accountability Incentive Block Grants Program Bulletin [On-line]. Available: http://www.ncjrs.org/html/ojjdp/jiabg_blltn_03_1_00/6.html.

Sample Qualified Service Organization Agreement and Consent Form

XYZ Service Center ("the Center") and

(Name of the alcohol/drug program)

(the "Program") hereby enter into an agreement whereby the Center agrees to provide:

_____

(Nature of services to be provided to the program)

Furthermore, the Center:

(1) Acknowledges that in receiving, transmitting, transporting, storing, processing, or otherwise dealing with any information received from the Program identifying or otherwise relating to the patients in the Program ("protected information"), it is fully bound by the provisions of the Federal regulations governing the Confidentiality of Alcohol and Drug Abuse Patient Records, 42 CFR, Part 2; and the Health Insurance Portability and Accountability Act (HIPAA), 45 CFR, Parts 142, 160, 162, and 164, and may not use or disclose the information except as permitted or required by this Agreement or by law;

(2) Agrees to resist any efforts in judicial proceedings to obtain access to the protected information except as expressly provided for in the regulations governing the Confidentiality of Alcohol and Drug Abuse Patient Records, 42 CFR, Part 2.

(3) Agrees to use appropriate safeguards (define with more specificity) to prevent the unauthorized use or disclosure of the protected information;

(4) Agrees to report to the Program any use or disclosure of the protected information not provided for by this Agreement of which it becomes aware (insert negotiated time and manner terms);

(5) Agrees to ensure that any agent, including a subcontractor, to whom the Center provides the protected information received from the Program, or created or received by the Center on behalf of the Program, agrees to the same restrictions and conditions that apply through this agreement to the Center with respect to such information;*

(6) Agrees to provide access to the protected information at the request of the Program, or to an individual as directed by the Program, in order to meet the requirements of 45 CFR § 164.524, which provides patients with the right to access and copy their own protected information (insert negotiated time and manner terms);

(7) Agrees to make any amendments to the protected information as directed or agreed to by the Program pursuant to 45 CFR § 164.526 (insert negotiated time and manner terms);

(8) Agrees to make available its internal practices, books, and records, including policies and procedures, relating to the use and disclosure of protected information received from the Program, or created or received by the Center on behalf of the Program, to the Program or to the Secretary of the Department of Health and Human Services for purposes of the Secretary determining the Program's compliance with HIPAA (insert negotiated time and manner terms);

(9) Agrees to document disclosures of protected information, and information related to such disclosures, as would be required for the Program to respond to a request by an individual for an accounting of disclosures in accordance with 45 CFR § 164.528 (insert negotiated time and manner terms);*

(10) Agrees to provide the Program or an individual information in accordance with paragraph (9) of this agreement to permit the Program to respond to a request by an individual for an accounting of disclosures in accordance with 45 CFR § 164.528 (insert negotiated time and manner terms);

Termination

(1) The program may terminate this agreement if it determines that the Center has violated any material term;

(2) Upon termination of this agreement for any reason, the Center shall return or destroy all protected information received from the Program, or created or received by the Center on behalf of the Program. This provision shall apply to protected information that is in the possession of subcontractors or agents of the Center. The Center shall retain no copies of the protected information.

(3) In the event that the Center determines that returning or destroying the protected information is infeasible, the Center shall notify the Program of the conditions that make return or destruction infeasible (insert negotiated time and manner terms).

Upon notification that the return or destruction of the protected information is infeasible, the Center shall extend the protections of this Agreement to such protected information and limit further uses and disclosures of the information to those purposes that make the return or destruction infeasible, as long as the Center maintains the information.

Executed this _____ day of _____, 20 ___.

_____ _____

President
XYZ Service Center

Program Director

Sample Consent Form

Consent for the Release of Con"dential Information

I, _____, authorize _____
(Name of patient) (Name or general designation of program making disclosure)

to disclose to _____
(Name of person or organization to which disclosure is to be made)

the follo wing information:_____
(Nature and amount of information to be disclosed; as limited as possible)

e purpose of the disclosure authorized in this is to:

_____
(Purpose of disclosure, as speci"c as possible)

_____

I understand that my substance use disorder treatment records are protected under the Federal regulations governing Con"dentiality and Drug Abuse Patient Records, 42 CFR, Part 2, and the Health Insurance Portability and Accountability Act of 1996, 45 CFR, Parts 160 and 164, and cannot be disclosed without my written consent unless otherwise provided for by the regulations. I also understand that I may revoke this consent in writing at any time except to the extent that action has been taken in reliance on it, and that in any event this consent expires automatically as follows:

_____
(Speci"cation of the date, event, or condition upon which this consent expires)

_____

I understand that generally (*insert name of program*) may not condition my treatment on whether I sign a consent form, but that in certain limited circumstances I may be denied treatment if I do not sign a consent form.

_____ _____
Date Signature of Patient

_____
Signature of parent, guardian, or uthorized representative where required.

Adapted from Legal Action Center. (n.d.). Sample consent - basic [On-line]. Available: http://www.lac.org/doc_library/lac/publications/Consent-Basic.pdf.

7KLV SDJH LV LQWHQWLRQDOO\ OHIW EODQ

7KLV SDJH LV LQWHQWLRQDOO\ OHIW EODQ